Newcastle University

A novel approach for holistic environmental assessment of ships

Mr. Martin Gibson, Dr Alan Murphy &
Dr Kayvan Pazouki

Shipping and the Environment Conference 2017

24-25th October 2017

CLINSH

email: m.t.gibson@newcastle.ac.uk

Lloyd's Register

Presentation Summary

- Aim and objectives
- Research background and question
- Assessment framework
- Method overview
- Initial results
- Conclusions and next steps



Aim and Objectives

Development of a framework and methodology for assessing the environmental impacts of ships using a holistic approach

- Overview of existing environmental initiatives in shipping
- Proposal of framework for ship environmental impact assessment
- Develop methodology for scoring impacts and prioritising environmental hazards
- Prioritise environmental hazards using proposed method

Research background and question

- Analysis of existing environmental initiatives in shipping
  - 74 initiatives identified
  - Categorised
  - Analysed: scope; weightings; and scoring range
- Limitations with existing initiatives identified
  - Applicability to ship type/location
  - Not ship specific
  - Rationale of scoring methods unclear
  - Indicator bias
  - Lack of ambition
  - Narrow scope

Alternative framework based on ship specific impacts

Framework


SCOPE
CONCEPTUAL MODEL
SOURCE
ENVIRONMENTAL INTERACTION
HAZARD
PATHWAY
ENVIRONMENTAL IMPACT
RECEPTOR
Spatial extent of impact, Temporal extent of impact, Visual impact
Impact Severity
Air, Earth System, Land, Water, Biota
Impact Groups
Legal requirement
Hazard Severity
Interacts with environment during normal shipping operation
Hazard Likelihood
Hazard Significance
Ship interactions with the environment

Newcastle University

Environmental interactions and hazards

INTERACTION	HAZARD
Discharges to sea	Oil
	Sewage
	Grey water
	Antifouling paint
	Invasive species transfer
	Marine litter
Emissions to air	GHG's (CO_2, Methane, N_2O, Halocarbons)
	SO_x
	NO_x
	Particulate
	VOC's
Anthropogenic Noise	Underwater noise
	Noise in port areas
Land	Waste (disposal)
	Resource depletion
Physical	Collisions with large aquatic life

References: (Andersson et al., 2016; IMO and the Environment, 2011; Talley, 2003)

Example

Newcastle University

Source-Pathway-Receptor example

ENVIRONMENTAL INTERACTION	HAZARD	SOURCE	PATHWAY	RECEPTOR	IMPACT
Emission to Air	CO_2	Fuel	Combustion	Atmosphere	Climate change Ocean acidification Disruption to carbon cycle
	Methane	LNG Fuel	Slippage due to incomplete combustion	Atmosphere	Climate change
	Nitrous oxide	Fuel	Combustion at low temp	Atmosphere	Climate change
	Halocarbons	Refrigerants	Leakage	Atmosphere	Climate change Ozone depletion
	SO_X	Fuel	Combustion	Atmosphere	Negative radiative forcing Acid rain Dry deposition
	NO_X	Fuel	Combustion (high temperature & low RPM)	Atmosphere	Marine eutrophication Ocean acidification Acid rain Low level ozone Secondary particulate formation Negative radiative forcing
	Particulate	Fuel; oil; components	Combustion; material wear	Atmosphere	Human inhalation (respiratory; lungs, heart) Negative radiative forcing Positive radiative forcing Cloud formation Decrease snow/ice albedo Acid rain
	VOC's	Crude oil; solvents	Evaporation; burning of marine fuel	Atmosphere	Human health - carcinogen Climate change Low level ozone

Newcastle University

Severity definitions

SEVERITY DEFINITIONS (to determine impact magnitude)		
Spatial extent	Global	Impacts the global environment e.g. a change in global atmospheric conditions
	Regional	Impacts the environment at a continental and/or national level
	Local	Impacts on the environment at a port or bay level
	Individual	Impacts which effect individual structures or organisms
	Negligible	No spatial impact on the environment
Temporal extent	Permanent	An impact with permanent or near permanent effects (i.e. > 100 years)
	Temporary	An impact with temporary effects (i.e. < 100 years)
	Immediate	An impact with immediate effects
	Negligible	An impact with no effects
Visual impact	Yes	Impacts are visible to naked eye
	No	Impacts cannot be seen
Legal requirement	Mandatory International	Hazard is covered by IMO mandatory international legislation
	International (not in force)	Hazard covered by IMO international legislation that has yet to enter force legally
	Optional International	Hazard is covered by IMO legislation classified as optional
	Regional	Hazard is not covered by IMO legislation but is covered by national or regional legislation (e.g. EU)
	No legislation	Hazard is not covered by legislation

Newcastle University

Impact severity indicators

	IMPACT SEVERITY			HAZARD SEVERITY
IMPACT LEVEL	**Spatial extent**	**Temporal extent**	**Visual impact**	**Legal Requirement**
5	Global	Permanent		Mandatory International
4	Regional			International (not in force)
3	Local	Temporary	Yes	Optional international
2	Individual			Regional
1	Negligible	Negligible	No	No legislation

Combining impacts into groups

HAZARD	IMPACTS	IMPACT MAGNITUDE						IMPACT SEVERITY
		SPATIAL	Score	TEMPORAL	Score	VISUAL	Score	
NO_X	Eutrophication	Regional	4	Temporary	3	No	1	12
	Ocean acidification	Regional	4	Temporary	3	No	1	12
	Acid rain formation	Local	3	Temporary	3	Yes	3	27
	Low level ozone formation	Local	3	Temporary	3	No	1	9
	Secondary particulate formation	Regional	4	Temporary	3	No	1	12
	Negative radiative forcing (cooling)	Regional	4	Temporary	3	No	1	12

HAZARD	IMPACTS	IMPACT GROUP				
		AIR	EARTH SYSTEM	LAND	WATER	BIOTA
NO_X	Eutrophication	×	×	×	✓	✓
	Ocean acidification	×	×	×	✓	✓
	Acid rain formation	✓	×	✓	✓	✓
	Low level ozone formation	✓	×	×	×	×
	Secondary particulate formation	✓	×	×	×	×
	Negative radiative forcing (cooling)	✓	✓	×	×	×

Determining hazard severity

HAZARD	IMPACT GROUP	IMPACT MAGNITUDE PER HAZARD						IMPACT GROUP SEVERITY	TOTAL SEVERITY
		SPATIAL	Score	TEMPORAL	Score	VISUAL	Score		
	AIR	Regional	4	Temporary	3	No	1	12	
	EARTH SYSTEM	Regional	4	Temporary	3	No	1	12	
	LAND	Local	3	Temporary	3	Yes	3	27	123
	WATER	Regional	4	Temporary	3	No	1	12	
NO_x	BIOTA	Regional	4	Permanent	5	Yes	3	60	



Example

Likelihood of hazard



Example

				LIKELIHOOD	
HAZARD	SOURCE	PATHWAY	RECEPTOR	ROUTINE INTERACTION WITH THE ENVIRONMENT	Score
CO_2	Fuel	Combustion	Atmosphere	Yes	5
Methane	LNG Fuel	Slippage due to incomplete combustion	Atmosphere	Yes	5
Nitrous oxide	Fuel	Combustion at low temp	Atmosphere	Yes	5
Halocarbons	Refrigerants	leakage	Atmosphere	No	1

Method - Calculations

Hazard significance = hazard severity x likelihood

Hazard severity = (Σ impact group severity) x legal requirement

Impact group severity = spatial x temporal x visual

Prioritising hazards - results

Discharges to sea
Emissions to air
Anthropogenic Noise
Land
Physical



Example

HAZARD	SCORE	RANKING
GHG's	3625	1
NO_x	1845	2
Particulate	1620	3
SO_x	1485	4
Waste (disposal)	1035	5
Resource depletion	615	6
Oil	390	7
Invasive species	300	8
VOC's	222	9
Antifouling paint	150	10
Noise (Underwater)	150	10
Sewage	135	12
Noise (ports)	60	13
Grey water	54	14
Marine litter	54	14
Collisions	16	16

Conclusions

- Development of method for prioritising hazards based on ship impacts
- Initial results consistent with industry expectation (LR consultation)

NEXT STEPS

- Further development of numerical indicators required to refine rankings
- Development of methodology required for analysis of ‘case study’ vessels
- Data collection for case study vessels to commence

Thank You

Mr. Martin Gibson, Dr Alan Murphy &
Dr Kayvan Pazouki

Shipping and the Environment Conference 2017

24-25th October 2017

email: m.t.gibson@newcastle.ac.uk